




Monterey Bay Aquarium Seafood Watch

Draft Assessment for Review April 2025

Octopus



© Scandinavian Fishing Yearbook

Species: Common octopus (*Octopus americanus* (*O. vulgaris*))
Mexican four-eyed octopus (*Octopus maya*)

Location: Mexico

Gear: Trolling lines

Type: Wild Caught

Author: Seafood Watch Consulting Researcher

Published:

Report ID: 28271

Assessed using [Seafood Watch Fisheries Standard v3](#)

Table of Contents

Table of Contents	2
About Monterey Bay Aquarium Seafood Watch	3
Seafood Watch Ratings	4
Guiding Principles	5
Final Ratings	8
Summary	9
Introduction	11
Assessments	18
Criterion 1: Impacts on the Species Under Assessment	18
Criterion 1 Summary	19
Criterion 1 Assessment	19
Criterion 2: Impacts on Other Species	37
Criterion 2 Summary	38
Criterion 2 Assessment	40
Criterion 3: Management Effectiveness	43
Criterion 3 Summary	44
Criterion 3 Assessment	44
Criterion 4: Impacts on the Habitat and Ecosystem	52
Criterion 4 Summary	53
Criterion 4 Assessment	53
Acknowledgements	57
References	58
Appendix A: Review Schedule	64

Draft for Review

About Monterey Bay Aquarium Seafood Watch

The mission of the Monterey Bay Aquarium is to inspire conservation of the ocean and enable a future where the ocean flourishes and people thrive in a just and equitable world. To do this, the Aquarium is focused on creating extraordinary experiences that inspire awe and wonder, championing science-based solutions, and connecting people across the planet to protect and restore the ocean. We know that healthy ocean ecosystems are critical to enabling life on Earth to exist, and that our very survival depends on them. As such, our conservation objectives are to mobilize climate action, improve the sustainability of global fisheries and aquaculture, reduce sources of plastic pollution, and restore and protect ocean wildlife and ecosystems.

The aquarium is focused on improving the sustainability of fisheries and aquaculture given the role seafood plays in providing essential nutrition for 3 billion people globally, and in supporting hundreds of millions of livelihoods. Approximately 180 million metric tons of wild and farmed seafood is harvested each year (excluding seaweeds). Unfortunately, not all current harvest practices are sustainable and poorly managed fisheries and aquaculture pose the greatest immediate threat to the health of the ocean and the economic survival and food security of billions of people.

The Seafood Watch program was started 25 years ago as a small exhibit in the Monterey Bay Aquarium highlighting better fishing practices and grew into one of the leading sources of information on seafood sustainability, harnessing the power of consumer choice to mobilize change. The program's comprehensive open-source information and public outreach raises awareness about global sustainability issues, identifies areas for improvement, recognizes and rewards best practices and empowers individuals and businesses to make informed decisions when purchasing seafood.

We define sustainable seafood as seafood from sources, whether fished or farmed, that can maintain or increase production without jeopardizing the structure and function of affected ecosystems, minimize harmful environmental impacts, assure good and fair working conditions, and support livelihoods and economic benefits throughout the entire supply chain. As one aspect of this vision, Seafood Watch has developed trusted, rigorous standards for assessing the environmental impacts of fishing and aquaculture practices worldwide. Built on a solid foundation of science and collaboration, our standards reflect our guiding principles for defining environmental sustainability in seafood.

Seafood Watch Ratings

The Seafood Watch Standard for Fisheries is used to produce assessments for wild-capture fisheries resulting in a Seafood Watch rating of green, yellow, or red. Seafood Watch uses the assessment criteria to determine a final numerical score as well as numerical subscores and colors for each criterion. These scores are translated to a final Seafood Watch color rating according to the methodology described in the table below. The table also describes how Seafood Watch defines each of these categories. The narrative descriptions of each Seafood Watch rating, and the guiding principles listed below, compose the framework on which the criteria are based.

Green	Final Score >3.2, and either criterion 1 or criterion 3 (or both) is green, and no red criteria, and no critical scores	Wild-caught and farm-raised seafood rated green are environmentally sustainable, well managed and caught or farmed in ways that cause little or no harm to habitats or other wildlife. These operations align with all of our guiding principles.
Yellow	Final score >2.2, and no more than one red criterion, and no critical scores, and does not meet the criteria for green (above)	Wild-caught and farm-raised seafood rated yellow cannot be considered fully environmentally sustainable at this time. They align with most of our guiding principles, but there is either one conservation concern needing substantial improvement, or there is significant uncertainty associated with the impacts of the fishery or aquaculture operations.
Red	Final Score \leq 2.2, or two or more Red Criteria, or one or more Critical scores.	Wild-caught and farm-raised seafood rated Red are caught or farmed in ways that have a high risk of causing significant harm to the environment. They do not align with our guiding principles and are considered environmentally unsustainable due to either a critical conservation concern, or multiple areas where improvement is needed.

Disclaimer: All Seafood Watch fishery assessments are reviewed for accuracy by external experts in ecology, fisheries science, and aquaculture. Scientific review does not constitute an endorsement of the Seafood Watch program or its ratings on the part of the reviewing scientists. Seafood Watch is solely responsible for the conclusions reached in this assessment.

Recommended Citation: Seafood Watch (2025) [Octopus](#) Monterey Bay Aquarium

Guiding Principles

Monterey Bay Aquarium defines sustainable seafood as seafood from sources, whether fished or farmed, that can maintain or increase production without jeopardizing the structure and function of affected ecosystems, minimize harmful environmental impacts, assure good and fair working conditions, and support livelihoods and economic benefits throughout the entire supply chain.

As one aspect of this vision, Seafood Watch has developed trusted, rigorous standards for assessing the environmental impacts of fishing and aquaculture practices worldwide. Environmentally sustainable wild capture fisheries:

1. Follow the principles of ecosystem-based fisheries management

The fishery is managed to ensure the integrity of the entire ecosystem, rather than solely focusing on maintenance of single species stock productivity. To the extent allowed by the current state of the science, ecological interactions affected by the fishery are understood and protected, and the structure and function of the ecosystem is maintained.

2. Ensure all affected stocks¹ are healthy and abundant

Abundance, size, sex, age and genetic structure of the main species affected by the fishery (not limited to target species) is maintained at levels that do not impair recruitment or long-term productivity of the stocks or fulfillment of their role in the ecosystem and food web.

Abundance of the main species affected by the fishery should be at, above, or fluctuating around levels that allow for the long-term production of maximum sustainable yield. Higher abundances are necessary in the case of forage species, in order to allow the species to fulfill its ecological role.

3. Fish all affected stocks at sustainable levels

Fishing mortality for the main species affected by the fishery should be appropriate given current abundance and inherent resilience to fishing while accounting for scientific uncertainty, management uncertainty, and non-fishery impacts such as habitat degradation.

¹“Affected” stocks include all stocks affected by the fishery, no matter whether target or bycatch, or whether they are ultimately retained or discarded.

The cumulative fishing mortality experienced by affected species must be at or below the level that produces maximum sustainable yield for single-species fisheries on typical species that are at target levels.

Fishing mortality may need to be lower than the level that produces maximum sustainable yield in certain cases such as forage species, multispecies fisheries, highly vulnerable species, or fisheries with high uncertainty.

For species that are depleted below target levels, fishing mortality must be at or below a level that allows the species to recover to its target abundance.

4. Minimize bycatch

Seafood Watch defines bycatch as all fisheries-related mortality or injury other than the retained catch. Examples include discards, endangered or threatened species catch, pre-catch mortality and ghost fishing. All discards, including those released alive, are considered bycatch unless there is valid scientific evidence of high post-release survival and there is no documented evidence of negative impacts at the population level.

The fishery optimizes the utilization of marine and freshwater resources by minimizing post-harvest loss and by efficiently using marine and freshwater resources as bait.

5. Have no more than a negligible impact on any threatened, endangered or protected species

The fishery avoids catch of any threatened, endangered or protected (ETP) species. If any ETP species are inadvertently caught, the fishery ensures and can demonstrate that it has no more than a negligible impact on these populations.

6. Are managed to sustain the long-term productivity of all affected species

Management should be appropriate for the inherent resilience of affected marine and freshwater life and should incorporate data sufficient to assess the affected species and manage fishing mortality to ensure little risk of depletion. Measures should be implemented and enforced to ensure that fishery mortality does not threaten the long term productivity or ecological role of any species in the future.

The management strategy has a high chance of preventing declines in stock productivity by taking into account the level of uncertainty, other impacts on the stock, and the potential for increased pressure in the future.

The management strategy effectively prevents negative population impacts on bycatch species, particularly species of concern.

7. Avoid negative impacts on the structure, function or associated biota of aquatic habitats where fishing occurs

The fishery does not adversely affect the physical structure of the seafloor or associated biological communities.

If high-impact gears (e.g. trawls, dredges) are used, vulnerable seafloor habitats (e.g. corals, seamounts) are not fished, and potential damage to the seafloor is mitigated through substantial spatial protection, gear modifications and/or other highly effective methods.

8. Maintain the trophic role of all aquatic life

All stocks are maintained at levels that allow them to fulfill their ecological role and to maintain a functioning ecosystem and food web, as informed by the best available science.

9. Do not result in harmful ecological changes such as reduction of dependent predator populations, trophic cascades, or phase shifts

Fishing activities must not result in harmful changes such as depletion of dependent predators, trophic cascades, or phase shifts.

This may require fishing certain species (e.g., forage species) well below maximum sustainable yield and maintaining populations of these species well above the biomass that produces maximum sustainable yield.

10. Ensure that any enhancement activities and fishing activities on enhanced stocks do not negatively affect the diversity, abundance, productivity, or genetic integrity of wild stocks

Any enhancement activities are conducted at levels that do not negatively affect wild stocks by reducing diversity, abundance or genetic integrity.

Management of fisheries targeting enhanced stocks ensures that there are no negative impacts on the wild stocks, in line with the guiding principles described above, as a result of the fisheries.

Enhancement activities do not negatively affect the ecosystem through density dependent competition or any other means, as informed by the best available science.

Final Ratings

Ratings Details	C 1 Target Species	C 2 Other Species	C 3 Managem ent	C 4 Habitat	Rating
Common octopus Mexico - Gulf of Mexico - Trolling lines - 1,900 mt	4.284	2.644	2.000	3.464	Yellow (2.976)
Mexican four-eyed octopus Mexico - Gulf of Mexico - Trolling lines - 23,000 mt	2.644	4.284	2.000	3.464	Yellow (2.976)

Draft for Review

Summary

The octopus' fishery is one of the most important in volume and value in the Gulf of Mexico. Producers in the Campeche Bank target the two main species, the Mexican four-eyed octopus (*Octopus maya*) and the common octopus (*Octopus americanus*).

This report assesses the octopus fisheries in the states of Campeche (composed mainly of *O. maya*) and Yucatan (*O. maya*, ~92%, and 3% *O. americanus*) (DOF 2024) targeted using baited trolling lines (known as "jimbas").

Until recently, abundance assessments were conducted by managers only for the Mexican four-eyed octopus. The assessments were used to estimate annual harvest quotas for the species. These evaluations were based on a dynamic biomass model (DOF 2014) that estimated the status against the MSY. However, in recent years, (Roa-Ureta 2021) recommended using the total latent productivity (TLP) as a more appropriate reference point to assess the stock status.

In 2024, managers assessed both species using the total latent productivity (TLP) (DOF b 2024). They concluded that neither species showed signs of overfishing (DOF 2024), which differed from the previous evaluation in 2022 (DOF 2022), where the fishery was considered deteriorated (DOF 2022). Due to conflicting results, a PSA analysis complemented the scoring for both species, and species abundance was deemed low concern for *O. americanus* and moderate for *O. maya*. Finally, fishing mortality was deemed a moderate concern, since levels of *F* are currently unknown.

Octopus trolling lines are highly selective, and no interactions with other species have been reported. Similarly, recent information regarding using protected species as bait has been released, so overall, the fishery does not raise concerns regarding bycatch.

Concerning management, a federal fishery management plan and a Mexican Official Norm are in place. These management tools contain specifications on the minimum size limit, the annual quota for red octopus, and a no-fishing season from 16 December 16 to July 31. These regulations have been partially effective, considering landings have been relatively steady, even though the red octopus quota (based on the MSY approach) was surpassed for several years. It is also important to consider that researchers have found misreporting to be an issue, and the current status of the species needs to be monitored closely. In addition, although efforts to improve enforcement and compliance have been promoted by authorities (Gobierno de Mexico 2019) and independent initiatives (Fishery Progress 2024), reports on significant use of illegal gear, noncompliance, and inadequate enforcement actions (Pradilla, A. 2019) (El Pais 2023) (InSight Crime 2021) (Oceana 2024) led the score of this subfactor as ineffective.

Finally, the gear used in the octopus fisheries in the Yucatan Peninsula has minimal impact on the benthic habitat due to its contact with the bottom. Still, some spatial regulations are in place to keep those impacts to a minimum. The ecosystem impacts from the trolling lines fishery are considered low.

Overall, the fisheries in Campeche and Yucatan that target common octopus (*O. americanus*)

and red octopus (*O. maya*) are rated yellow.

Draft for Review

Introduction

Scope of the analysis and ensuing rating

This Seafood Watch assessment provides recommendations for the red octopus (*Octopus maya*) and common octopus (*Octopus americanus*) fisheries in the Campeche Bank, Mexico. The species are targeted by two fleets (a small-scale fleet and a “semi” industrial fleet comprised of 10 mt of storage vessels that operate as motherships of small-scale ships known as “alijos”) in the states of Yucatan and Campeche (DOF 2024). Both fleets use baited lines (or jimbas) that drift on the bottom of the sea from their vessels or alijos (Figure 2).

Species Overview

Red octopus (*O. maya*) is an endemic benthic species living in shallow waters of the continental shelf of the Yucatan (INAPESCA 2014). Its known distribution extends from the waters adjacent to Isla del Carmen in Campeche to Isla Mujeres in Quintana Roo (DOF 2014). One single stock is exploited based on genetic studies. This species has been extensively studied in the region due to its high value and importance for the regional economy.

The common octopus, a member of a species complex (*O. vulgaris*) composed of six species (Maloney et al 2023) was recently renamed as *O. americanus*, and its presence under this name was confirmed along the Yucatan continental shelf by the genetic results of (Avendaño et al. 2020). It is exploited in the Yucatan Peninsula and Veracruz (DOF 2014) (see Figure 1).




Figure 1: Distribution of the different species of the *O. vulgaris* complex, including *O. americanus* (*O. maya* and *O. insularis* are included) (Avendaño et al. 2020)

Once they hatch, red octopus juveniles behave like adults, capturing live prey and remaining

hidden when inactive (Santamaría et al. 2023). Reproduction is continuous, with greater intensity towards the end of each year. Fecundity fluctuates between 1,500 and 2,000 eggs per female (Silva et al 2002). Because almost all mature females and those who have spawned are less likely to feed, the fishing method employed in the Yucatan Peninsula is considered a very conservative method, since it could capture more males than females (DOF 2014) (see figure 2). Overall, the red octopus has rapid growth and a short life cycle (Solis-Ramirez and Chavez 1986). The maximum age has been estimated at around 12 months and 18 months (Solis-Ramirez and Chavez 1986)(Arreguin-Sanchez 1992). (Angeles-Gonzalez et al. 2018) analyzed the spatio-temporal variability of the reproductive maturation for red octopus and reported that high temperatures (close to and above 27 °C) significantly impact the distribution and spawning of the species. In addition, it has been reported that the reproductive strategy of red octopus females ensures that an important fraction of eggs hatches, combined with two annual recruitment events, contribute to the population biomass (Duarte et al 2018).



Figure 2: Fishing vessels with Jimbas (bamboo sticks) with baited trolling lines

On the other hand, the common octopus is a cosmopolitan species distributed in tropical and

subtropical waters worldwide. It can be found from the sea surface to a depth of 150 m. It lives in waters with temperatures between 6 °C and 33 °C, being more common between 10 °C and 30 °C (INAPESCA b 2014).

The common octopus reaches its sexual maturity at 8 to 10 months when they measure 125 mm of mantle length (ML) (Santos-Valencia et al. 2018) Solis-Ramirez et al. 1997}. However, it has been suggested that the size of sexual maturity differs between sexes. In the Gulf of Mexico, Jimenez-Badillo (2008) found the same pattern described in the Mediterranean, where the minimum size of sexual maturity in males was 248 g and in females 612 g (Jimenez-Badillo 2008). On average, most males reach sexual maturity at 110 mm ML with 700 g total weight, while the females reach it at 140 mm ML with 1,400 g total weight (Silva et al 2002). The spawning period is continuous throughout the year in tropical and subtropical waters. A female specimen can deposit between 100,000 and 500,000 eggs during its short lifetime (Perez et al 2007). This high level of fecundity coincides with the reports in the Canary Islands, which found that the real fecundity oscillated between 31 and 106 eggs spawned per gram of female body weight (Hernandez-Garcia et al. 2002). Growth estimates for the Gulf of Mexico suggest slower growth than in the Mediterranean and a maximum age of 2.2 years (Diaz-Alvarez and Jimenez-Badillo 2009).

Production Statistics

In the state of Yucatan, octopus captures have varied over time. From 1998 to 2000, landings remained between 14,000 and 15,000 tons (t); from 2001 to 2003, less than 10,000 t per year was recorded. During the 2004 to 2010 period, there was an alternation between low (2005, 2008) and high (2004, 2006, 2009) capture levels. In 2006 the historical maximum of the series was obtained with around 20,000 t, which was very close during the 2009 season. Based on the official data report, from 2006 to 2021, landings were relatively stable, around the 20,000 t starting in 2009 to 2013 and above 24,000 t from 2014 to 2019, with a decrease in 2020 (mostly driven by the Covid pandemic) to reach almost 30,000 t in 2021 (see images below) (CONAPESCA 2021). On the other hand, Campeche landings increased steadily from 1998 to 2001, reaching the historical maximum of 8,107 t in 2001. Since 2002, annual catches have remained above 5,000 t, except for the 2008 season, when the catch decreased to 3,946 t. After that year, landings remained around the 20,000 and 25,000 t range (CONAPESCA 2024).




Figure 3: Yucatan and Campeche octopus historical landings (DOF 2024)

According to CONAPESCA data, production in Campeche is dominated by red octopuses. In Yucatan, ~60% is red octopuses and 40% is other octopus species (see figures below).




Figure 4: Yucatan octopus' species proportions in 2024. Built using data from CONAPESCA 2024




Figure 5: Campeche octopus species catch composition in 2024. Built using data from CONAPESCA

Importance to the US/North American market

In 2022, Mexico was still among the most important octopus producers in the world, covering around 1.6% of the total export volume in the world (OEC 2024). That year, exports were destined mainly to the USA (43%), Italy (16%), Colombia (13%), Portugal (9%), Greece (7%), and Japan (6%) (OEC 2024). According to NOAA Foreign trade data, between 2021 and 2023, an average of 2,293 metric tons of octopus were imported from Mexico; most of the production from the region is reaching the export market (El Pais 2023).

Table 1

Product	2022	2023	2024
Octopus dried/salt/brine	1,803,789	1,707,051	728,724
Octopus frozen	254,968	234,126	167,488
Octopus NSPF/prepared/preserved	77,300	107,848	18,632

Table 2

Product	2021	2022	2023
OCTOPUS DRIED/SALTED/BRINE	348.7	255.0	234.1
OCTOPUS FROZEN	2340.9	1803.8	1707.1
OCTOPUS LIVE/FRESH	4.8	0.0	0.0
OCTOPUS NSPF PREPARED/PRESERVED	1.2	77.3	107.8

Common and market names

Red octopus (*Octopus maya*) is also known as octopus maya in Mexico or Mexican four-eye octopus (Fish Source 2024). *Octopus americanus* is known as octopus common, or “patón” in Mexico, and common octopus in the US (FDA 2018).

Primary product forms

Octopus landed in the Yucatan Peninsula is available in the local, national, and international markets (Coronado et al. 2020). Whole octopus, fresh and iced, is marketed domestically. Exported octopus is whole frozen and packed in 20 kg boxes. An organism’s size classifies octopuses. Octopus trade is carried out by permit holders, retailers, middlemen in the Yucatan Peninsula, and national seafood traders (Coronado et al. 2020).

Production volume notes

Landings are part of the 2024 official landing information (CONAPESCA 2024) for *O. maya* and *O. americanus* in the states of Campeche and Yucatan.

Summary

The fisheries in Campeche and Yucatan that target common octopus (*O. americanus*), and red octopus (*O. maya*) are rated yellow.

Eco-Certification information

The octopus fishery is engaged in two Fishery Improvement Projects (FIP). Engagement in a FIP does not affect the Seafood Watch score per se because we base our assessments on the

current situation. However, data and information generated by these improvement projects can be used in our assessments.

Draft for Review

Assessments

This section assesses the sustainability of the fishery(s) relative to the Seafood Watch Standard for Fisheries, available at www.seafoodwatch.org. The specific standard used is referenced on the title page of all Seafood Watch assessments.

Criterion 1: Impacts on the Species Under Assessment

This criterion evaluates the impact of fishing mortality on the species, given its current abundance. When abundance is unknown, abundance is scored based on the species' inherent vulnerability, which is calculated using a Productivity-Susceptibility Analysis. The final Criterion 1 score is determined by taking the geometric mean of the abundance and fishing mortality scores. The Criterion 1 rating is determined as follows:

- Score >3.2 = **Green** or Low Concern
- Score >2.2 and ≤ 3.2 = **Yellow** or Moderate Concern
- Score ≤ 2.2 = **Red** or High Concern

Rating is Critical if Factor 1.3 (Fishing Mortality) is Critical.

Guiding principles

- *Ensure all affected stocks are healthy and abundant.*
- *Fish all affected stocks at sustainable level*

Criterion 1 Summary

Common octopus			
Region / Method	Abundance	Fishing Mortality	Score
Mexico - Gulf of Mexico - Trolling lines - 1,900 mt	3.670 Low Concern	5.000 Low Concern	Green (4.284)

Mexican four-eyed octopus			
Region / Method	Abundance	Fishing Mortality	Score
Mexico - Gulf of Mexico - Trolling lines - 23,000 mt	2.330 Moderate Concern	3.000 Moderate Concern	Yellow (2.644)

Criterion 1 Assessment

Scoring Guidelines

Factor 1.1 - Abundance

Goal: Stock abundance and size structure of native species is maintained at a level that does not impair recruitment or productivity.

- 5 (Very Low Concern) — Strong evidence exists that the population is above an appropriate target abundance level (given the species' ecological role), or near virgin biomass.
- 3.67 (Low Concern) — Population may be below target abundance level, but is at least 75% of the target level, OR data-limited assessments suggest population is healthy and species is not highly vulnerable.
- 2.33 (Moderate Concern) — Population is not overfished but may be below 75% of the target abundance level, OR abundance is unknown and the species is not highly vulnerable.
- 1 (High Concern) — Population is considered overfished/depleted, a species of concern, threatened or endangered, OR abundance is unknown and species is highly vulnerable.

Factor 1.2 - Fishing Mortality

Goal: Fishing mortality is appropriate for current state of the stock.

- 5 (Low Concern) — Probable (>50%) that fishing mortality from all sources is at or below a sustainable level, given the species ecological role, OR fishery does not target species and fishing mortality is low enough to not adversely affect its population.
- 3 (Moderate Concern) — Fishing mortality is fluctuating around sustainable levels, OR fishing mortality relative to a sustainable level is uncertain.
- 1 (High Concern) — Probable that fishing mortality from all source is above a sustainable

level.

Draft for Review

Common octopus (*Octopus americanus* (*O. vulgaris*))

1.1 Abundance

Gulf of Mexico - Trolling lines

Low Concern

In 2021, (Roa-Ureta 2021) released the results of a stock assessment using an alternative model on *O. americanus*, which was recently identified as a distinct species of the *O. vulgaris* complex (Avendaño et al. 2020). The stock assessment used an intra-annual generalized depletion (IAGD) and the Pella-Tomlinson models to estimate the species biomass and reference points (Roa-Ureta 2021), including the target latent productivity (TLP) as an alternative reference point instead of the “traditional” MSY. The TLP was proposed as a sustainable annual harvest rate for fluctuating stocks by (Poot-Salazar et al 2024).

Based on the results, (Roa-Ureta 2021) concluded that although the stock appeared to have a flat trajectory (see image in justification), there was evidence of fluctuations that occur under disturbances (e.g., catches higher than the historical average) (ibid). In particular, several start-of-season biomass estimates from IAGD were imprecise, and the stock tended to enter a fluctuating regime under relatively moderate fishing pressure (i.e., annual catches higher than 12 to 15 thousand t) (Roa-Ureta 2021). The author stated that MSY estimates were nearly four times higher than the reference point estimated based on latent productivity; however, despite the imprecise estimate of latent productivity, the reference point was much more conservative and considered the fluctuating nature of the stock (ibid).

Most recently, an assessment update was released (with data from the 2022 and 2023 seasons) and included the estimated status of the *O. americanus* using the MSY and the TLP as references in Kobe plots (see figures in justification). The graphs showed that when MSY is used as a reference point, $B < B_{MSY}$; however, when TLP is used, $B > B_{TLP}$.

(Roa-Ureta 2021)(Roa-Ureta 2024) concluded that the stock of *O. americanus* appears to be under-fished. However, whether the assessment underwent a robust scientific peer review is still being determined. Still, the conclusion aligns with the most recent National Fisheries Chart statement that reported the species has “development opportunities” (DOF 2022).

Finally, in 2024, managers released an update on the stock status using the (Roa-Ureta 2024) approach and the TLP as a reference point (see Figure 12 in justification). Managers concluded that the species biomass is above the reference point and the fishery has “a potential to develop more” (DOF b 2024).

The most recent results from the data-limited assessment completed by managers showed

that the species biomass is above the limit reference point appropriate for the species (TLP) (Poot-Salazar et al 2024), and since there are no conflicting indicators, this factor is scored as a “low concern”.

Supplementary Information




Figure 6: Start of the season biomass for *O. americanus* (*O. vulgaris*) (Roa-Ureta 2021)




Figure 7: Status of *O. americanus* using the MSY (top) and TLP (bottom) as reference points (Roa-Ureta 2024)




Figure 8: Biomass dynamics (green dots and black lines), landings (blue bars) and sustainable catches (Total Latent Productivity) for common octopus in the Yucatan Peninsula. Areas around the red lines are standard error bands (DOF b 2024)

1.2 Fishing Mortality

Gulf of Mexico - Trolling lines

Low Concern

Fishing mortality reference points have not been set for the common octopus in the Gulf of Mexico. According to the 2022 National Fisheries Chart, the fishery is at its maximum and no increase in effort is recommended (DOF 2018)(DOF 2022). During the most recent update of the fishery profile, it was reported that one medium scale vessel and more than 1,400 small-scale boats were actively fishing in Campeche, while ~441 medium-scale and more than 3,200 small-scale vessels were also actively fishing in Yucatan (DOF 2022).

The 2021 stock assessment estimated fishing mortality values for the species from 2000 to 2019 (see images below) (Roa-Ureta 2021). The stock assessment defines the instantaneous exploitation rate as the ratio between fishing mortality and natural mortality. The fishing mortality was lower than natural mortality in 2000, 2002, 2003, 2005 to 2010, 2012 to 2016, 2018 and 2019. In 2001 and 2004 the fishing mortality and natural mortality were very close (Roa-Ureta 2021) (see image in justification). It was concluded that the

instantaneous exploitation rate for *O. americanus* in Yucatan was within the sustainable regime (Roa-Ureta 2021).

Additionally, based on official reports, landings for *Octopus* spp. in Yucatan showed an increasing trend from 2008 (2,436 t) to a maximum of 15,525 t in 2018. In 2020, production was significantly reduced, due to the COVID-19 pandemic, and in 2021, landings reached more than 12000 (CONAPESCA 2021).

Considering that landings have remained relatively stable, that managers reported the species has potential for growth (DOF 2022), and the analysis developed by (Roa-Ureta 2021) reported similar findings, this factor is a "low concern" for common octopus.

Supplementary Information

Draft for Review




Figure 9: Fishing mortality (red line) and natural mortality (green line) rates estimated from IAGD models for the *O. americanus* stock in the Yucatan Peninsula, Mexico. (Roa-Ureta 2021)

Mexican four-eyed octopus (*Octopus maya*)

1.1 Abundance

Gulf of Mexico - Trolling lines

Moderate Concern

Scientific surveys and catch data have been used to estimate the abundance of red octopus in the past. Managers have in place annual monitoring programs to assess and suggest annual quotas (DOF 2014). Studies have been completed recently to understand the species' status better.

In 2019, (Avendaño et al. 2019) used four different methodologies to assess the biomass of the species in deeper areas (between 30 and 60 meters depth) of the Campeche Bank. The authors concluded that the fraction of the population that occurs between that range of depth consisted mainly of adult organisms so that it could contribute significantly to the recruitment of the entire population and recommended that fishing in these depths should be avoided (Avendaño et al. 2019).

(Roa-Ureta 2021) completed a stock assessment for *O. maya* for the Gulf of Mexico and the Caribbean, using a detailed database of fishing trips that included quantities of total catch and fishing effort from 2000 to 2019. The author used an alternative model with the intra-annual generalized depletion model (IAGD) to estimate the biomass at the start of the seasons and the Pella-Tomlinson production model to calculate MSY and the total latent productivity (Roa-Ureta 2021). TLP was proposed as a sustainable annual harvest rate for fluctuating stocks by (Poot-Salazar et al 2024).

(Roa-Ureta 2021) estimates for K , MSY, B_{MSY} , and latent productivity (see table below) were calculated following that approach and concluded that latent productivity was 50% higher than the mean annual catch over the studied period (2000 to 2019). Latent productivity was very close to the current harvest rate, which was only exceeded in 2018 (ibid).

Table 3

Parameter	Maximum likelihood estimate	Standard error	Lower bound	Upper bound
$r (y-1)$	3.382	2.072		
p	1.876	0.474		
$k (t)$	292,044	66,168		

MSY (t)	225,092	45,179	134,734	315,450
B_{MSY}	142,433	45,094	52,245	232,621
Latent productivity	18,012	7,592	2,827	33,196

In 2024, an assessment update that includes data from the 2022 and 2023 seasons was released, and the estimated status of the *O. maya* was included using the MSY and the TLP as reference points in Kobe plots (see figures in justification) (Roa-Ureta 2024). The graphs showed that when MSY is used as a reference point, $B < B_{MSY}$; however, when TLP is used, $B > B_{TLP}$.

Finally, and as part of the 2024 update of the fishery's profile within the National Fisheries Chart, managers released the results of the stock evaluation for *O. maya* using the TLP - suggested by (Roa-Ureta 2024)- as a reference point to estimate the status of the species. Managers reported that the species was exploited to its maximum sustainable yield, with biomass above the TLP reference point (see figure 10) (DOF 2024), an improvement from the previous update (DOF 2022) where the species was reported to be deteriorated and $B_{2019} < B_{MSY}$ (Figure 8 in justification).

Since there are data-limited assessments (Poot-Salazar et al 2024)(Roa-Ureta 2024) that provide conflicting conclusions (when using MSY and TLP), and the species is not highly vulnerable, abundance is scored as a "moderate concern".

Supplementary Information




Figure 10: Kobe plot for *O. maya* in the Yucatan Peninsula (DOF 2022)




Figure 11: Start-of-season biomass population dynamics of *O. maya* in the Yucatan Peninsula. Dots of varying size and grayness are estimated from IAGD models (Roa-Ureta 2021)




Figure 12: *O. maya* status using MSY (top) and TLP (bottom) estimated by (Roa-




Figure 13: Biomass dynamics (green dots and black lines), landings (blue bars) and sustainable catch rates (TLP) of the red octopus (*Octopus maya*) in the Yucatan Peninsula (DOF b 2024) The red arrow indicates the year in which changes in biomass dynamics parameters occur at: the symmetry of the production function (p) in *O. maya* (red areas around the red line are standard error bands).

Table 4

Table 5		
Productivity Attribute	Relevant Information	Score (1 = low risk, 2 = medium risk, 3 = high risk)
Average age at maturity	< 1 year (Rosas et al. 2014)	1
Average maximum age	12 months (Rosas et al. 2014)	1
Fecundity	>100,000 oocytes (Avila-Poleda et al. 2016)	1
Average maximum size (fish only)	N/A	
Average size at maturity (fish only)	N/A	
Reproductive strategy	Small hatchlings with pelagic paralarvae. (Lima et al. 2014)	2
Trophic level	~3.74	3
Density dependence (invertebrates only)	Unlikely (highly mobile species) (Salas et al. 2009)	2
Habitat Quality	N/A	
Total Productivity (average)	1.60	

Table 6		
Table 7		
Susceptibility Attribute	Relevant Information	Score (1 = low risk, 2 = medium risk, 3 = high risk)

<p>Areal overlap (Considers all fisheries)</p>	<p>The species distributes along the Yucatan Peninsula in the Gulf of Mexico (considered an endemic species) (Sealife Base 2024). Considering the importance of the species and the available data regarding fishing grounds (DOF 2022) >30% of the species is fished.</p>	<p>3</p>
<p>Vertical overlap (Considers all fisheries)</p>	<p>According to (Hochberg and Camacho-García 2009) the species ranges from 0 to 100 m. According to managers, the small-scale fleets that target octopuses in Campeche and Yucatan reach up to 37 m in depth (DOF 2014), and for these reasons, this factor scores as medium risk.</p>	<p>2</p>
<p>Selectivity of fishery (Specific to fishery under assessment)</p>	<p>Mexican federal legislation established a minimum legal size of 11 cm of mantle length (DOF 2016). For this reason, it is believed that all organisms within the landings are adults; therefore, individuals smaller than size at maturity are not regularly caught.</p>	<p>2</p>
<p>Post-capture mortality (Specific to fishery under assessment)</p>	<p>Octopus maya is targeted using live bait that guarantees that undersized organisms can be returned alive (DOF 2014). However, there is no evidence of a post-capture release; for this reason, this factor scores as medium risk.</p>	<p>2</p>

PSA score for <i>O. maya</i> is calculated as follows:
--

$$\text{Vulnerability (V)} = \sqrt{P^2 + S^2}$$

$$V = \sqrt{1.66^2 + 1.58^2}$$

$$V = 2.50$$

1.2 Fishing Mortality

Gulf of Mexico - Trolling lines

Moderate Concern

For *Octopus maya*, managers set quotas typically set based on the maximum sustainable yield (MSY) (DOF 2022). However, there are no set values related to F_{MSY} . Within the 2022 update on the fishery's profile, managers reported that, as of 2019 (the last year assessed by managers), fishing mortality was above the F_{MSY} (see Kobe plot in justification). This has also been the case for the 2016 to 2018 period (DOF 2022).

During the most recent update to the fishery's profile, managers stated that the *O. maya* is exploited within the maximum sustainable yield, however, there were no estimates related to F (DOF 2024). According to official landings (see introduction), during the most recent season, landings did not reach the authorized quota (DOF b 2024). Based on the fact that currently levels of F are unknown, this factor is scored as moderate concern.

Supplementary Information




Figure 10: Kobe plot for *O. maya* in the Yucatan Peninsula (DOF 2022)

Criterion 2: Impacts on Other Species

All main retained and bycatch species in the fishery are evaluated under Criterion 2. Seafood Watch defines bycatch as all fisheries-related mortality or injury to species other than the retained catch. Examples include discards, endangered or threatened species catch, and ghost fishing. Species are evaluated using the same guidelines as in Criterion 1. When information on other species caught in the fishery is unavailable, the fishery's potential impacts on other species is scored according to the Unknown Bycatch Matrices, which are based on a synthesis of peer-reviewed literature and expert opinion on the bycatch impacts of each gear type. The fishery is also scored for the amount of non-retained catch (discards) and bait use relative to the retained catch. To determine the final Criterion 2 score, the score for the lowest scoring retained/bycatch species is multiplied by the discard/bait score. The Criterion 2 rating is determined as follows:

- Score >3.2 = **Green** or Low Concern
- Score >2.2 and ≤ 3.2 = **Yellow** or Moderate Concern
- Score ≤ 2.2 = **Red** or High Concern

Rating is Critical if Factor 2.3 (Fishing Mortality) is Critical

Guiding principles

- Ensure all affected stocks are healthy and abundant.
- Fish all affected stocks at sustainable level.
- Minimize bycatch.

Criterion 2 Summary

Criterion 2 score(s) overview

This table(s) provides an overview of the Criterion 2 subscore, discards+bait modifier, and final Criterion 2 score for each fishery. A separate table is provided for each species/stock that we want an overall rating for.

Common octopus			
Region / Method	Sub Score	Discard Rate/Landings	Score
Mexico - Gulf of Mexico - Trolling lines - 1,900 mt	2.644	1.000: < 100%	Yellow (2.644)

Mexican four-eyed octopus			
Region / Method	Sub Score	Discard Rate/Landings	Score
Mexico - Gulf of Mexico - Trolling lines - 23,000 mt	4.284	1.000: < 100%	Green (4.284)

Criterion 2 main assessed species/stocks table(s)

This table(s) provides a list of all species/stocks included in this assessment for each 'fishery' (as defined by a region/method combination). The text following this table(s) provides an explanation of the reasons the listed species were selected for inclusion in the assessment.

Gulf of Mexico - Trolling lines			
Sub Score: 2.644	Discard Rate: 1.000		Score: 2.644
Species	Abundance	Fishing Mortality	Score
Mexican four-eyed octopus	2.330: Moderate Concern	3.000: Moderate Concern	Yellow (2.644)
Common octopus	3.670: Low Concern	5.000: Low Concern	Green (4.284)

Octopus fishing is done exclusively during the day, using a fishing technique known as "gareteo", which consists of leaving the boat drifting and dragging lines where bait is held (see image below)(DOF 2014). According to the Octopus Management plan, the fishing method is practically exclusive for capturing the target species, and confirmed that there is no incidental fishing during octopus capture operations (ibid). The most common species used in the red octopus fishery are mainly crabs (*Callinectes spp.*) and spider crab (*Libinia dubia*)(DOF 2014).




Figure 16: Fishing vessels with Jimbas (bamboo sticks) with baited trolling lines (DOF 2014)

For that reason, no bycatch species are included in the recommendation. Because the octopus species in this report are sometimes caught together, each is included as a Criterion 2 species for the other octopus species.

Although it has been reported that one of the species used as bait is the endangered horseshoe crab (Smith et al 2017), a species protected under Mexican federal legislation (NOM-059), the level of impact on horseshoe crabs from this fishery is unclear, so it is not included in the assessment.

Criterion 2 Assessment

Scoring Guidelines

Factor 2.1 - Abundance

(same as Factor 1.1 above)

Factor 2.2 - Fishing Mortality

(same as Factor 1.2 above)

Factor 2.3 - Modifying Factor: Discards and Bait Use

Goal: Fishery optimizes the utilization of marine and freshwater resources by minimizing post-harvest loss. For fisheries that use bait, bait is used efficiently.

Scoring Guidelines: The discard rate is the sum of all dead discards (i.e. non-retained catch) plus bait use divided by the total retained catch.

Ratio of bait + discards/landings	Factor 2.3 score
<100%	1
>=100	0.75

Draft for Review

2.3 Discard Rate/Landings

Gulf of Mexico - Trolling lines

< 100%

The fishing technique used in the octopus fishery is very selective, and no incidental catch is recorded. According to the management plan, several crab species are the main groups used as bait (DOF 2014).

One of the fishery improvement projects (FIP), has a monitoring program that reports on production and collects relevant information regarding bait use. Data recorded by the FIP includes the 2021, 2022, and 2023 seasons. Results indicate that the fishery uses three main species as bait, with mangrove crab (*Ucides cordatus*) as the most important (averaging ~70%). The spider crab (*Libinia dubia*)(~18%), and the Chesapeake blue crab (*Callinectes sapidus*)(~12%) complete the list of the species mostly used as bait (Barajas et al. 2022) (Ibarra-García et al. 2023) (Barajas et al. 2024).

One of the bait species mentioned in the literature was the horseshoe crab (*Limulus polyphemus*), a protected species under Mexican legislation, within the NOM-059. Although recent information to corroborate this was not available, according to Smith et al. {2017}, there were increasing reports of small-scale poaching of horseshoe crab adults used solely as an alternative to commercial bait species in the artisanal octopus fishery of Campeche and Yucatan. In addition, in 2023 (Munguia-Vega et al. 2023) conducted a study using DNA barcoding to identify species used as bait from 12 localities during 2022. The authors reported nine crustacean species. Of these, the mangrove crab was the most frequent (61%), followed by spider crab (14%) and Caribbean spiny lobster (*Panulirus argus*, 8%). The report confirmed that the American horseshoe crab (*Limulus polyphemus*), was absent in the samples. During the last reported seasons, the amount of bait used relative to total landings was significantly below the catches (see table below). For this reason, this factor is scored as <100.

Table 8

Season	2021	2022	2023
# of logbooks registered	1,883	1,340	3,429
Octopus production (kg)	1,407,700	986,523	788,687
Bait used (kg)	86,330	29,129	52,473

Octopus species	<i>O. maya</i> and <i>O. americanus</i>	<i>O. maya</i> and <i>O. americanus</i>	<i>O. maya</i> and <i>O. americanus</i>
Bait species used	<i>U. cordatus</i> , <i>L. dubia</i> and <i>C. sapidus</i>	<i>U. cordatus</i> , <i>L. dubia</i> and <i>C. sapidus</i>	<i>U. cordatus</i> , <i>L. dubia</i>

Draft for Review

Criterion 3: Management Effectiveness

Five factors are evaluated in Criterion 3: Management Strategy and Implementation, Bycatch Strategy, Scientific Research/Monitoring, Enforcement of Regulations, and Inclusion of Stakeholders. Each is scored as either 'highly effective', 'moderately effective', 'ineffective,' or 'critical'. The final Criterion 3 score is determined as follows:

- 5 (Very Low Concern) — Meets the standards of 'highly effective' for all five factors considered.
- 4 (Low Concern) — Meets the standards of 'highly effective' for 'management strategy and implementation' and at least 'moderately effective' for all other factors.
- 3 (Moderate Concern) — Meets the standards for at least 'moderately effective' for all five factors.
- 2 (High Concern) — At a minimum, meets standards for 'moderately effective' for Management Strategy and Implementation and Bycatch Strategy, but at least one other factor is rated 'ineffective.'
- 1 (Very High Concern) — Management Strategy and Implementation and/or Bycatch Management are 'ineffective.'
- 0 (Critical) — Management Strategy and Implementation is 'critical'.

The Criterion 3 rating is determined as follows:

- Score >3.2 = **Green** or Low Concern
- Score >2.2 and ≤3.2 = **Yellow** or Moderate Concern
- Score ≤2.2 = **Red** or High Concern

Rating is Critical if Management Strategy and Implementation is Critical.

Guiding principle

- The fishery is managed to sustain the long-term productivity of all impacted species.

Five factors are evaluated in Criterion 3: Management Strategy and Implementation, Bycatch Strategy, Scientific Research/Monitoring, Enforcement of Regulations, and Inclusion of Stakeholders. Each is scored as either 'highly effective', 'moderately effective', 'ineffective,' or 'critical'. The final Criterion 3 score is determined as follows:

Criterion 3 Summary

Fishery	Management Strategy And Implementation	Bycatch Strategy	Scientific Research And Monitoring	Enforcement Of Management Regulations	Stakeholder Inclusion	Score
Gulf of Mexico - Trolling lines	Moderately Effective	Highly effective	Moderately Effective	Ineffective	Highly effective	Red (2.000)

Criterion 3 Assessment

Scoring Guidelines

Factor 3.1 - Management Strategy and Implementation

Considerations: What type of management measures are in place? Are there appropriate management goals, and is there evidence that management goals are being met? Do managers follow scientific advice? To achieve a highly effective rating, there must be appropriately defined management goals, precautionary policies that are based on scientific advice, and evidence that the measures in place have been successful at maintaining/rebuilding species.

Factor 3.2 - Bycatch Strategy

Considerations: What type of management strategy/measures are in place to reduce the impacts of the fishery on bycatch species and when applicable, to minimize ghost fishing? How successful are these management measures? To achieve a Highly Effective rating, the fishery must have no or low bycatch, or if there are bycatch or ghost fishing concerns, there must be effective measures in place to minimize impacts.

Factor 3.3 - Scientific Research and Monitoring

Considerations: How much and what types of data are collected to evaluate the fishery's impact on the species? Is there adequate monitoring of bycatch? To achieve a Highly Effective rating, regular, robust population assessments must be conducted for target or retained species, and an adequate bycatch data collection program must be in place to ensure bycatch management goals are met.

Factor 3.4 - Enforcement of Management Regulations

Considerations: Do fishermen comply with regulations, and how is this monitored? To achieve a Highly Effective rating, there must be regular enforcement of regulations and verification of compliance.

Factor 3.5 - Stakeholder Inclusion

Considerations: Are stakeholders involved/included in the decision-making process? Stakeholders are individuals/groups/organizations that have an interest in the fishery or that may

be affected by the management of the fishery (e.g., fishermen, conservation groups, etc.). A Highly Effective rating is given if the management process is transparent, if high participation by all stakeholders is encouraged, and if there a mechanism to effectively address user conflicts.

Draft for Review

3.1 Management Strategy And Implementation

Gulf of Mexico - Trolling lines

Moderately Effective

The current management strategies for the octopus fishery are based mostly on red octopus data and scientific information. A limited amount of information (e.g., landing data) is collected for common octopus, and the recently identified variant of *O. insularis*. Access is controlled with the use of fishing licenses, and there is a minimum size limit of 110 cm, which applies to all octopus species in the Gulf of Mexico and Caribbean (DOF 2016). The exception is for *O. americanus* females within the Veracruz Reef National Park, where the minimum size limit is set at 140 mm; this measure aims to allow organisms to breed or reproduce before they are caught (DOF 2014)(DOF 2015)(DOF 2016). The federal Red and Common Octopus Fishery Management Plan (OFMP) includes an annual closed fishing season that runs from December to July (DOF 2014)(DOF 2016).

In addition, for *Octopus maya*, an estimated and suggested quota is determined based on adult abundance, recruitment, and growth data estimates (INAPESCA 2014) (INAPESCA b 2014) (Perez et al 2007). According to managers, these quotas are typically set based on the maximum sustainable yield (MSY), a level of fishing that allows the population to persist. The goal is to allow 50% of the population to escape the fishery and survive to reproduce (INAPESCA b 2014). Managers followed scientific recommendations, and quotas coincided with what was advised until 2014, 2020, and 2022 (DOF b 2024). As of 2001, the catch quota was assigned according to an annual abundance assessment and has varied between 10,000 t and 13,000 t, with 10,000 t (INAPESCA 2014) (figure below). We built a graph to present official landing data versus set limits to visualize production against the recommended quota (see image below). It is also important to consider that, according to (Suasnávar-Imán et al 2021), the decisions on the quota have not incorporated elements from population dynamics necessary for sustainable exploitation. Based on the available data, the quota seems to be surpassed several times (see table and figure in justification below). In addition, and no catch quotas are set for the other species.

Finally, in terms of the volume of landings, *O. maya* represented roughly 90% of the fishery's main targeted species. The current management plan (DOF 2014) as well as the Federal Official Norm (DOF 2015) are currently based on this species' biology, and the measures are expected to be effective. In recent years, the limit quota has been constantly surpassed, there is a need for increased precaution. As part of these efforts, a monitoring program has been in place that follows landings and updates are publicly available (FIP Pulpo Yucatan 2024); once the fishery reaches the quota, the fishery will close. However, it is unclear what effective mechanisms are in place to effectively close the fishery, and if these are effective. For these reasons, this factor is scored as a moderate concern.

Supplementary Information




Figure 14: *O. maya* landings and quota between 2006 and 2023 (CONAPESCA 2021) (Gobierno de Mexico 2024) (INAPESCA b 2014) (INAPESCA 2014)

3.2 Bycatch Strategy

Gulf of Mexico - Trolling lines

Highly effective

According to the information available, the trolling lines or *jimbos* have no interaction with other species and for that reason a bycatch strategy is unnecessary. This factor is scored as "highly effective."

3.3 Scientific Research And Monitoring

Gulf of Mexico - Trolling lines

Moderately Effective

The fishery considers the three octopus species as target species (DOF 2022). Both management instruments (the fishery management plan and the Mexican official norm (DOF 2014) (DOF 2015)) consider only two of the species within their scope (*O. maya* and *O. americanus*). In addition, the efforts to estimate biomass and assign quotas are only developed for *O. maya* (DOF 2022). These are calculated by managers at the Mexican Center for Research on Fisheries and Aquaculture (IMIPAS) who use CPUE data to fit a Schaefer surplus production model (Roa-Ureta 2021). The CPUE is generated from producers' landing tickets.

Based on these efforts, managers at CONAPESCA recommended measures in line with a reduction of effort, an increase in monitoring, control, and enforcement, and ensuring the implementation of the management plan (DOF 2022). Still, it was concluded that for *O. americanus*, the fishery has the potential to develop even further within the 36 to 150 meters range of depth (DOF 2022). However, the document did not share details on how this conclusion was generated.

In the case of *O. maya*, some stock abundance and health data are collected and analyzed (DOF 2022), which is used to monitor and maintain the stock using appropriate data-limited assessment methods (Surplus production model) and management strategies. For *O. americanus* and *O. insularis*, minimal data are collected or analyzed (landings and potential use of CPUE), and management methods are similar to those applied for *O. maya*. Overall, management relies on one strategy requiring minimal monitoring (CPUE from landing tickets).

O. maya has historically been the main driver of the fishery, in terms of proportion of the catch (DOF 2014). Abundance estimates have been analyzed and quotas have been generated for the species. Recently, data on landings were used also to estimate the status of *O. maya* and *O. americanus* (Roa-Ureta 2024). It is essential to consider that details on *O. insularis* are minimal and could make the score ineffective. Considering that *O. maya* represents >70% of the catch this score remains as “moderately effective.”

3.4 Enforcement Of Management Regulations

Gulf of Mexico - Trolling lines

Ineffective

Participation in the fishery is controlled by permits that provide access to the species to the holders (DOF 2022). The enforcement of regulations is carried out by the CONAPESCA officials (via federal officers), who are in charge of patrolling the coastline of more than 11,000 km. However, in recent years, the number of federal inspectors has decreased, from approximately 210 in 2012 to 161 in 2022 (Oceana 2024), which may be an insufficient number of agents for this amount of coastline (IMCO 2012) (Oceana 2024).

In 2020, the Mexican organization Causa Natura released the “Effectiveness Index for the Fishery Inspection and Enforcement in Mexico” (Pescando Datos 2020). The authors measure the effectiveness of fisheries surveillance in each state of the Mexican Republic using information on the level of actions deployed, the resources invested in inspection and surveillance, and how these were reflected in the results obtained. The authors built an index that included information from the National Aquaculture and Fisheries Commission (CONAPESCA). The report concluded that for Yucatán, the index was 0.37 in the scale of 0 to 1 (Pescando Datos 2020). In 2021, the state reported a “slight” improvement in the index value (0.552) (Pescando datos 2021) although still reporting improvement in areas related to better enforcement actions (Pescando datos 2021) (see image in justification).

In a similar approach, Oceana Mexico released a report analyzing official information related to five elements of enforcement actions in Mexico. These elements were:

- the number of federal enforcement agents
- number of enforcement actions (in water and land)

- amount of fish products and gears retained as part of enforcement actions
- the level of coverage of the VMS system (for industrial fleet)
- the number of sanctions and infractions applied.

Based on the analysis of the information on those five components, the report concluded that in the best of cases, there has been a stagnation of adequate enforcement actions by the authorities. In contrast, in the worst case, the results represent an apparent disregard for the State to fulfill its duties (Oceana 2024).

In addition, news outlets have recently released special reports that focused on the octopus fishery and the lack of effective enforcement, the increase in illegal activities that include using unauthorized gear (diving or cement bricks), and the consensus from researchers, activists, and fisher representatives, that better enforcement and compliance are needed (InSight Crime 2021) (El Pais 2023) (Balam, L. 2023)

This information contrasts with the official announcements for a collaboration agreement formalized between the Navy and the Ministry of Agriculture and Rural Development in 2019 (Gobierno de Mexico b 2019). The program aimed to strengthen inspection and surveillance tasks related to poaching fishing, which has since increased the surveillance capacity at sea and the coast.

Finally, one of the two Fishery Improvement Projects designed and launched a community sanction mechanism for non-compliance (Fishery Progress 2024). The FIP managers reported that the mechanism is considered to have a dissuasive and practical effect on FIP participants. Also, the existence of a monitoring, control, and surveillance mechanism in place during the 2022 fishing season, which was audited and resulted in a "Satisfactory compliance" as proof of effective compliance (Ecomorphoses 2023) (Fishery Progress 2024). In addition, FIP participants have held workshops in coordination with government agencies, academic institutions, and the fishing sector, to reinforce knowledge about octopus fishing management tools (Gamboa-Álvarez-Barajas-Girón b 2024) (Gamboa-Álvarez & Barajas-Girón 2024), although currently specific actions in place to improve enforcement are not in place.

In addition, these results only apply to the FIP participants, which, according to the FIP profile, covers ~14% of the fishery production (landings), but only around 3.7% of small-scale and 8.3% of mid-scale vessels (171 and 37, respectively) according to the FIP vessel list, compared to the active fleet reported by managers (4,707 small-scale and 442 mid-scale vessels) (Fishery Progress 2024).

Overall, enforcement actions are in place to promote management benefits. However, based on the reports and existing evidence, this might be inadequate for the scale of the fishery, and there seems to be constant poor compliance. For these reasons, this factor is scored as "ineffective."

Supplementary Information



Figure 15: Enforcement Effectiveness Index values for Mexican states in 2021 (including Yucatan) (Pescando datos 2021)

3.5 Stakeholder Inclusion

Gulf of Mexico - Trolling lines

Highly effective

The Comite Estatal Sistema Producto (CESP) was created for the octopus fishery in Yucatan (Por Esto! 2019). The CESP is a legal body recognized by managers that coordinates planning, communication, and builds agreements between the various participants in the fishery and supply chain (CONAPESCA 2016). Legal entities (e.g., civil associations) and other fishery stakeholders actively participate in the CESP meetings and create strategic planning instruments (e.g., master programs, guiding plans, and multi-year planning sheets) to improve their competitiveness and expand market participation (CONAPESCA 2016). One example of this collaboration was the management plan (DOF 2014) and the update to the official norm (DOF 2016) developed in coordination with scientists and other stakeholders, including the CESP.

In addition, and as part of one of the active FIPs, workshops were held with the fishing sector of Campeche and Yucatán to provide feedback on the Octopus Fisheries Management Plan (PMPP) and management measures. This workshop focused on reinforcing knowledge about the management tools in the Campeche Bank fishery, Mexico,

as well as developing strategies to implement actions to ensure their compliance (Gamboa-Álvarez & Barajas-Girón 2024).

Due to the successful involvement of diverse stakeholder groups in fishery management, this factor is scored as “highly effective.”

Draft for Review

Criterion 4: Impacts on the Habitat and Ecosystem

This Criterion assesses the impact of the fishery on seafloor habitats, and increases that base score if there are measures in place to mitigate any impacts. The fishery's overall impact on the ecosystem and food web and the use of ecosystem-based fisheries management (EBFM) principles is also evaluated. Ecosystem Based Fisheries Management aims to consider the interconnections among species and all natural and human stressors on the environment. The final score is the geometric mean of the impact of fishing gear on habitat score (factor 4.1 + factor 4.2) and the Ecosystem Based Fishery Management score. The Criterion 4 rating is determined as follows:

- Score >3.2 = **Green** or Low Concern
- Score >2.2 and ≤3.2 = **Yellow** or Moderate Concern
- Score ≤2.2 = **Red** or High Concern

Guiding principles

- Avoid negative impacts on the structure, function or associated biota of marine habitats where fishing occurs.
- Maintain the trophic role of all aquatic life.
- Do not result in harmful ecological changes such as reduction of dependent predator populations, trophic cascades, or phase shifts.
- Ensure that any enhancement activities and fishing activities on enhanced stocks do not negatively affect the diversity, abundance, productivity, or genetic integrity of wild stocks.
- Follow the principles of ecosystem-based fisheries management.

Rating cannot be Critical for Criterion 4.

Criterion 4 Summary

Fishery	Impact of Fishing Gear on the Habitat/Substrate	Modifying Factor: Mitigation of Gear Impacts	Ecosystem-based Fisheries Management	Score
Gulf of Mexico - Trolling lines	Score: 4	Score: 0	Moderate Concern	Green (3.464)

Criterion 4 Assessment

Scoring Guidelines

Factor 4.1 - Physical Impact of Fishing Gear on the Habitat/Substrate

Goal: The fishery does not adversely impact the physical structure of the ocean habitat, seafloor or associated biological communities.

- 5 - Fishing gear does not contact the bottom
- 4 - Vertical line gear
- 3 - Gears that contacts the bottom, but is not dragged along the bottom (e.g. gillnet, bottom longline, trap) and is not fished on sensitive habitats. Or bottom seine on resilient mud/sand habitats. Or midwater trawl that is known to contact bottom occasionally. Or purse seine known to commonly contact the bottom.
- 2 - Bottom dragging gears (dredge, trawl) fished on resilient mud/sand habitats. Or gillnet, trap, or bottom longline fished on sensitive boulder or coral reef habitat. Or bottom seine except on mud/sand. Or there is known trampling of coral reef habitat.
- 1 - Hydraulic clam dredge. Or dredge or trawl gear fished on moderately sensitive habitats (e.g., cobble or boulder)
- 0 - Dredge or trawl fished on biogenic habitat, (e.g., deep-sea corals, eelgrass and maerl)

Note: When multiple habitat types are commonly encountered, and/or the habitat classification is uncertain, the score will be based on the most sensitive, plausible habitat type.

Factor 4.2 - Modifying Factor: Mitigation of Gear Impacts

Goal: Damage to the seafloor is mitigated through protection of sensitive or vulnerable seafloor habitats, and limits on the spatial footprint of fishing on fishing effort.

- +1 —>50% of the habitat is protected from fishing with the gear type. Or fishing intensity is very low/limited and for trawled fisheries, expansion of fishery's footprint is prohibited. Or gear is specifically modified to reduce damage to seafloor and modifications have been shown to be effective at reducing damage. Or there is an effective combination of 'moderate' mitigation measures.
- +0.5 —At least 20% of all representative habitats are protected from fishing with the gear type and for trawl fisheries, expansion of the fishery's footprint is prohibited. Or gear modification measures or other measures are in place to limit fishing effort, fishing

intensity, and spatial footprint of damage caused from fishing that are expected to be effective.

- *0 — No effective measures are in place to limit gear impacts on habitats or not applicable because gear used is benign and received a score of 5 in factor 4.1*

Factor 4.3 - Ecosystem-Based Fisheries Management

Goal: All stocks are maintained at levels that allow them to fulfill their ecological role and to maintain a functioning ecosystem and food web. Fishing activities should not seriously reduce ecosystem services provided by any retained species or result in harmful changes such as trophic cascades, phase shifts or reduction of genetic diversity. Even non-native species should be considered with respect to ecosystem impacts. If a fishery is managed in order to eradicate a non-native, the potential impacts of that strategy on native species in the ecosystem should be considered and rated below.

- *5 — Policies that have been shown to be effective are in place to protect species' ecological roles and ecosystem functioning (e.g. catch limits that ensure species' abundance is maintained at sufficient levels to provide food to predators) and effective spatial management is used to protect spawning and foraging areas, and prevent localized depletion. Or it has been scientifically demonstrated that fishing practices do not have negative ecological effects.*
- *4 — Policies are in place to protect species' ecological roles and ecosystem functioning but have not proven to be effective and at least some spatial management is used.*
- *3 — Policies are not in place to protect species' ecological roles and ecosystem functioning but detrimental food web impacts are not likely or policies in place may not be sufficient to protect species' ecological roles and ecosystem functioning.*
- *2 — Policies are not in place to protect species' ecological roles and ecosystem functioning and the likelihood of detrimental food impacts are likely (e.g. trophic cascades, alternate stable states, etc.), but conclusive scientific evidence is not available for this fishery.*
- *1 — Scientifically demonstrated trophic cascades, alternate stable states or other detrimental food web impact are resulting from this fishery.*

4.1 Impact of Fishing Gear on the Habitat/Substrate

Gulf of Mexico - Trolling lines

Score: 4

The octopus fishery in the Gulf of Mexico uses trolling lines or *jimbass* that have minimal impact on the bottom. The bait normally comes in contact with the seafloor (DOF 2016) (DOF 2014). Fishing lines are pulled in by hand, so the gear has no mechanical elements. Considering that the different species of octopus prefer either seagrass, rocky, and coral reef habitats, fishing may occur over these areas (DOF 2014); however, because the gear does not damage bottom habitats, this factor is scored as 4 based on the SFW Fisheries Standard.

4.2 Modifying Factor: Mitigation of Gear Impacts

Gulf of Mexico - Trolling lines

Score: 0

In the Gulf of Mexico and the Mexican Caribbean, there are protected areas (under different categories, e.g., National Park, or Biosphere Reserves) with differing levels of protection; in some instances, certain fishing gears are banned, or fishing may be prohibited. However, no specific restrictions are in place for the octopus fishery beyond those included in the NOM and the Management Plan. For these reasons, no mitigation credit is awarded.

4.3 Ecosystem-based Fisheries Management

Gulf of Mexico - Trolling lines

Moderate Concern

Octopus species that are actively targeted have high trophic levels, and have a primarily carnivorous diet (e.g., crabs, mollusks and fish). All three species are also an important food source for other species in the ecosystem, like groupers and snappers. (Arreguin-Sanchez et al. 2000) (DOF 2016).

Commercially important spiny lobster and red grouper share habitat with octopus; these species are all connected through the food web. Octopuses are an important prey item for red grouper, and both red grouper and octopus are predators of spiny lobster. Fishing and management regulations for any of these species will likely have some effect on the others (Lasseter 2006) (Arreguin-Sanchez et al. 2000). Unfortunately, the roles of the octopus species in the food web are not well studied. Also, there has been no assessment of the fishery's effects on resources commonly used for bait (e.g., crabs). Since there is no

management or assessment of ecosystem impacts, but octopuses are not considered to be playing an exceptionally large role in the ecosystem, this factor is rated as “moderate concern”.

Draft for Review

Acknowledgements

Scientific review does not constitute an endorsement of the Seafood Watch® program, or its seafood recommendations, on the part of the reviewing scientists. Seafood Watch® is solely responsible for the conclusions reached in this report.

Draft for Review

References

- Angeles-Gonzalez, L., Calva, R., Santos-Valencia, J., Avila-Poveda, O., Olivares, A., Diaz, F., and Rosas, C. 2017. Temperature modulates spatio-temporal variability of the functional reproductive maturation of *Octopus maya* (Cephalopoda) on the shelf of the Yucatán Peninsula, Mexico. *Journal of Molluscan Studies* (2017) 83: 280-288
- Arreguin-Sanchez, F., 1992. Growth and seasonal recruitment of the octopus (*Octopus maya*) fishery from the Campeche Bank. *NAGA ICLARM Quarterly*. https://digitalarchive.worldfishcenter.org/bitstream/handle/20.500.12348/2985/na_2869.pdf?sequence=1&isAllowed=y
- Arreguin-Sanchez, F., Soliz-Ramirez M.J., Gonzalez de la Rosa M.E. 2000. Population dynamics and stock assessment for *Octopus maya* (Cephalopoda: Octopodidae) fishery in the Campeche Bank, Gulf of Mexico. *Rev. biol. trop.* Vol. 48, no. 2-3, p. 323-331.
- Avendaño O, Velázquez– Abunader I, Fernández–Jardón C, Ángeles– González LE, Hernández-Flores A, Guerra Á (2019). Biomass and distribution of the red octopus (*Octopus maya*) in the north-east of the Campeche Bank. *Journal of the Marine Biological Association of the United Kingdom* 99, 1317–1323. <https://doi.org/10.1017/S0025315419000419>
- Avendaño, O., Roura, A., Cedillo-Robles, C., González, A., Rodriguez, R., Velazquez-Abunader, I. and Guerra, A. 2020. *Octopus americanus*: a cryptic species of the *O. vulgaris* species complex redescribed from the Caribbean.
- Avila-Poveda, O.H., Koueta, N., Villalobos, F., Santos-Valencia, J., and Rosas, C. 2016. Reproductive traits of *Octopus maya* (Cephalopoda: Octopoda) with implications for fisheries management. *Molluscan Research*. 36. 29-44. [10.1080/13235818.2015.1072912](https://doi.org/10.1080/13235818.2015.1072912).
- Balam, L. 2023. Pesca ilegal del pulpo en Yucatán evidencia protección 'de chocolate' a la especie. *Causa Natura Media*. <https://causanaturamedia.com/periodismo-cn/pesca-ilegal-del-pulpo-en-yucatan-evidencia-proteccion-de-chocolate-a-la-especie>
- Barajas-Girón, M.P., Rocha-Tejeda, L., López-Ercilla, I. 2022. Reporte ejecutivo de monitoreo pesquero piloto implementando PescaData en el FIP de pulpo de Yucatán, temporada 2021. *Comunidad y Biodiversidad, A.C.* 15 pp.
- Barajas-Girón, P., Rocha-Tejeda, L., López-Ercilla, I. 2024. Fishing monitoring by implementing PescaData, to determine octopus catches and the use of bait species in the octopus FIP in Yucatán, 2023 season. *Community and Biodiversity, A.C.* 21 pages.

CONAPESCA 2016. El Comité Sistema Producto, un detonante para la competitividad.

CONAPESCA 2021. Anuario Estadístico de Acuicultura y Pesca de la Comisión Nacional de Acuicultura y Pesca.

CONAPESCA. (2024). Prensa: Está México en tercer lugar en producción de pulpo a nivel mundial, de acuerdo a la última publicación de la FAO.

Coronado, E., Salas, S., Cepeda-González, M. F., & Chuenpagdee, R. (2020). Who's who in the value chain for the Mexican octopus fishery: Mapping the production chain. *Marine Policy*, 118, 104013.

Coronado, E., Salas, S., Torres-Irineo, E., & Chuenpagdee, R. (2020). Disentangling the complexity of small-scale fisheries in coastal communities through a typology approach: The case study of the Yucatan Peninsula, Mexico. *Regional Studies in Marine Science*, 36, 101312. <https://doi.org/10.1016/j.rsma.2020.101312>

Díaz -Alvarez, A.G. y L. Jimenez-Badillo. 2009. Edad y crecimiento de pulpo (*Octopus vulgaris*; Cuvier, 1797) en el Parque Nacional Sistema Arrecifal Veracruzano. 1er Simposium para el conocimiento de los recursos costeros del Sureste de México, Ciudad del Carmen, Campeche, México.

DOF 2014. ACUERDO por el que se da a conocer el Plan de Manejo Pesquero de pulpo (O. Maya y O. Vulgaris) del Golfo de México y Mar Caribe. <https://inapesca.gob.mx/portal/documentos/Planes-de-Manejo-Pesquero/Golfo/Plan-de-Manejo-Pesquero-de-Pulpo.pdf>

DOF 2015. NORMA Oficial Mexicana NOM-008-SAG/PESC-2015, Para ordenar el aprovechamiento de las especies de pulpo en las aguas de jurisdicción federal del Golfo de México y Mar Caribe.

DOF 2016. NORMA Oficial Mexicana NOM-008-SAG/PESC-2015, Para ordenar el aprovechamiento de las especies de pulpo en las aguas de jurisdicción federal del Golfo de México y Mar Caribe.

DOF 2018.

ACUERDO por el que se da a conocer la actualización de la Carta Nacional Pesquera. DOF: 11/06/2018

DOF 2022. ACUERDO mediante el cual se da a conocer la actualización de la Carta Nacional Pesquera.

DOF 2024 ACUERDO por el que se establece el volumen de captura permisible para el

aprovechamiento del recurso pulpo rojo (*Octopus maya*) en aguas de jurisdicción federal en los estados de Campeche y Yucatán para la temporada de pesca 2024.

DOF 2024 b. Acuerdo mediante el cual se da a conocer la actualización de la Carta Nacional Pesquera

Duarte, J. A., Hernandez-Flores, A., Salas, S., and Seijo, J. C. 2018. Is it sustainable fishing for *Octopus maya* Voss and Solis, 1996, during the breeding season using a bait-based fishing technique? *Fisheries Research* 199 (2018) 119-126.

Ecomorphoses 2023. Resultados de la implementación del protocolo de auditorías para el cumplimiento de la normativa pesquera en el FIP de pulpo Yucatán

El Pais 2023. La depredación del pulpo bebé: ¿el fin de la pesca de la esperanza en el sur de México? El pulpo maya da sustento a unos 20.000 pescadores en la Península de Yucatán. Se exporta principalmente a Europa. Pero su captura ilegal y la sobreexplotación ponen en riesgo la salud de los pescadores y los ecosistemas.

FDA 2018. The FDA Seafood List - Updates for 2018

FIP Pulpo Yucatan 2024. Avance de las capturas de Pulpo Rojo

Fish Source 2024. Mexican four-eyed octopus Yucatan Peninsula.
https://www.fishsource.org/stock_page/2288

Fishery Progress 2024. Mexico Yucatan octopus - drift rod and line (Alianza Por El Pulpo En Yucatan)

Gamboa-Álvarez & Barajas-Girón 2024. Reporte del tercer taller anual de la pesquería de pulpo en la península de Yucatán.

Gamboa-Álvarez-Barajas-Girón b 2024. Reporte del taller de seguimiento a las estrategias para el respeto de las herramientas de manejo de Pulpo (HMPP).

Gobierno de México 2019. Esperanza de los pescadores yucatecos: el pulpo. El 1º de agosto inició la temporada de captura de pulpo en Yucatán, y las expectativas son prometedoras para una comunidad de aproximadamente 15 mil pescadores, entre ribereños y tripulantes de los barcos de la flota mayor.

Gobierno de Mexico 2024. Otorgan Indicación Geográfica al pulpo Maya de la península de Yucatán, primera pesquería mexicana considerada un bien nacional: Agricultura.
Secretaría de Agricultura y Desarrollo Rural | 25 de agosto de 2024 | Comunicado.

Gobierno de Mexico b 2019. Fortalecen acciones SADER y SEMAR en beneficio del sistema

productivo pesquero y acuícola.

Secretaría de Agricultura y Desarrollo Rural | 11 de febrero de 2019 | Comunicado

Hernández-García, V & Hernández-López, J.L. and Castro, J. 2002. On the reproduction of *Octopus vulgaris* off the coast of the Canary Islands. *Fisheries Research*. 57. 197-203. 10.1016/S0165-7836(01)00341-1.

Hochberg, F.G. and Y.E. Camacho-García 2009 *Squids and octopuses*. p. 399-408 Wehrmann, I.S.; Cortés, J. 2009. *Marine biodiversity of Costa Rica, Central America*. Springer 538pp.

Ibarra-García, L, Barajas-Girón, M.P., Rocha-Tejeda, L., López-Ercilla, I. 2023. Fishing monitoring by implementing PescaData, and characterization of bait species in the Octopus FIP in Yucatan, 2022 season. *Community and Biodiversity*, A.C. 15 pages.

IMCO 2012. LA PESCA ILEGAL E IRREGULAR EN MÉXICO: UNA BARRERA A LA COMPETITIVIDAD

INAPESCA b 2014. *Sustentabilidad y Pesca Responsable en Mexico*. Evaluación y Manejo. Documento interno, Instituto Nacional de Pesca.

INAPESCA. 2014. Dictamen Técnico. Establecimiento de cuota de captura de pulpo (*Octopus maya*) para la temporada de pesca 2014. Documento interno. Instituto Nacional de Pesca.

InSight Crime 2021. *Tentacles of Illegal Octopus Trade Clutch Mexico Town*.

Jimeñez-Badillo, M.L., 2008. Management challenges of the small scale fishing communities in a protected reef system of Veracruz, Gulf of Mexico. *Journal of the Fisheries Management and Ecology*, 15: 19-26.

Lasseter, Ava. 2006. "Lobsters are like Gold': Perceptions of resource access and management in a Mexican spiny lobster fishery" M.A. Thesis. University of Florida, Gainesville, Florida, USA. 180 pp.

Lima, F., Leite, T., Haimovici, M., Nóbrega, M., and Lins-Oliveira, J. 2014. Population structure and reproductive dynamics of *Octopus insularis* (Cephalopoda: Octopodidae) in a coastal reef environment along northeastern Brazil. *Fisheries Research*. 152. 86–92. 10.1016/j.fishres.2013.08.009.

Maloney B, Ramos EA, Bennice CO, Young F, Magnasco MO. Genetic confirmation of *Octopus insularis* (Leite and Haimovici, 2008) in South Florida, United States using physical features and de novo genome assembly. *Front Physiol*. 2023 Jun 20;14:1162807. doi: 10.3389/fphys.2023.1162807. PMID: 37408588; PMCID: PMC10318165.

Munguia-Vega, Adrian & García, Laura & Barajas-Girón, Polo & Tejeda, Lorena & Lopez-Ercilla, Ines & Domínguez-Contreras, Jose. (2023). Genetic Identification of Bait to Support Sustainability of the Octopus Fishery from the State of Yucatan, Mexico. *Journal of Shellfish Research*. 42. 10.2983/035.042.0212.

Oceana 2024. PESCA ILEGAL EN MÉXICO: SOLUCIONES DESDE LA POLÍTICA PESQUERA

OEC 2024. Molluscs; octopus (*Octopus* spp), frozen.

Perez, M., R. Burgos, A.T. Wakida-Kusunoki, J. Santos, K. Cervera, J.C. Espinoza, J.C. Mena, E. Cob y M. Medina. 2006. Evaluacion de la poblacion de pulpo *Octopus maya* en la Peninsula de Yucatan. 2006. Informe de Investigacion. Instituto Nacional de Pesca. CRIP-Yucalpeten. Doc. Interno 11 p.

Pescando Datos 2020. Índice de efectividad en inspección y vigilancia pesquera

Pescando datos 2021.
Índice de Efectividad
en Inspección y Vigilancia.

Poot-Salazar A, Vela-zquez-Abunader I, Avendaño O, Barajas-Girón P, Rojas-González RI, Pensamiento-Villarauz S, et al. (2024) Sustainable fishing harvest rates for fluctuating fish and invertebrate stocks. *PLoS ONE* 19(9): e0307836.
<https://doi.org/10.1371/journal.pone.0307836>

Por Esto! 2019. Instalan primer Comité Consultivo del Pulpo.
<https://www.poresto.net/yucatan/2019/7/17/instalan-primer-comite-consultivo-del-pulpo.html>

Pradilla, A. 2019. Pesca ilegal 'agota' el pulpo, el mero y el pepino marino en Yucatán. *Animal Politico*. <https://www.animalpolitico.com/2019/09/pesca-ilegal-crisis-yucatan-pulpo-mero>

Roa-Ureta 2024. Desarrollo de diagramas de Kobe basados en la Productividad Latente Total y actualización a 2023 de evaluaciones de stock de *Octopus maya* y *O. americanus* capturados en Yucatán, México.

Roa-Ureta, R. 2021. Stock Assessment of *Octopus maya* and *O. americanus* in Mexican waters of the Gulf of Mexico and the Caribbean. August 11, 2021.

Rosas, C., Gallardo, P., Mascaró, M., Caamal-Monsreal, C., and, & Pascual, C. 2014. *Octopus maya*. *Cephalopod Culture*. 383-396. 10.1007/978-94-017-8648-5_20.

Salas, S., M.A. Cabrera, L. Palomo y E. Torres. 2009. Uso de Indicadores para Evaluar Medidas de Regulación en la Pesquería del Pulpo en Yucatán dada la Interacción de Flotas.

Santamaría A, Euán Ávila JI, Munguia-Rosas MA, Saldívar-Lucio R and Fraga J (2023) Environmental variability and governance: The fishery of Octopus maya in Yucatan, Mexico. Front. Mar. Sci. 10:1018728. doi:10.3389/fmars.2023.1018728” Excerpt From Environmental variability and governance: The fishery of Octopus maya in Yucatan, Mexico <https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2023.1018728/epub>

Santos-Valencia, Josefina & Can-González, Maricarmen & Velazquez-Abunader, Iván & Avendaño, Otilio. (2018). First estimates about reproductive aspects of common octopus (*Octopus vulgaris*) at northeast Yucatan Peninsula, Mexico. November 2018. Conference: Cephalopod International Advisory Council Conference 2018. Cephalopod Research Across Scales: From Molecules to Ecosystems At: St. Petersburg, Florida, USA

Sealife Base 2024. Octopus maya Voss & Solis Ramirez, 1966 Mexican four-eyed octopus.

Silva, L., Sobrino, I. & Ramos, F. 2002. Reproductive biology of the common octopus, *Octopus vulgaris* Cuvier, 1797 (Cephalopoda: Octopodidae) in the Gulf of Cadiz (SW Spain). Bulletin of Marine Science 71,837-850

Smith, D. R., Brockmann, H. J., Beekey, M. A., King, T. L., Millard, M. J., & Zaldivar-Rae, J. (2017). SCS Global Services Report Conservation status of the American horseshoe crab, (*Limulus polyphemus*): a regional assessment. Reviews in Fish Biology and Fisheries, 27(1), 135-175.

Solis-Ramirez M.J., Chavez E.A. 1986. Evaluacion y regimen optimode pesca de pulpo de la Peninsula de Yucatan, Mexico. Ann. Inst. Cienc. Mar Limnol. Univ. Nac. Aut. Mex. 13: 1–18.

Suasnívar-Imán, M., Rocha-Tejeda, L., Barajas-Girón, M.P. y López-Ercilla, I. 2021. Monitoreo de Plan de Manejo Pesquero de pulpo (*O. maya* y *O. vulgaris*) del Golfo de México y Mar Caribe. Comunidad y Biodiversidad, A.C. 29 pp.

Appendix A: Review Schedule

This report was reviewed in March 2025 and updated from version 3 of the Fisheries Standard to version 4. The changes are described as follows:

- *O. insularis* was removed from the recommendation, considering the limited proportion of their catch (0.1%) in the fishery
- C1 Abundance for *O. maya* and *O. americanus* upgraded to moderate concern based on a recent evaluation conducted by managers to assess the status of the stocks.
- C2 The removal of *O. insularis* from this criterion and the lack of bycatch associated with this fishery improved the scoring from a “red” to a “yellow” scoring for C2.
- The overall rating for the fishery improved from red to yellow for both species remaining.

Draft for Review